

4.3/4.4, Model Input Namelist Parameters, (online), available from <<http://www.atmet.com/> html/docs/documentation.shtml>, (accessed 2005-09-07).

High Resolution Wind Flow Simulation Using a Meso-scale Meteorological Model : Application to Wind Power Generation Plant over the Noma Cape, Kagoshima

Keiya YUMIMOTO*, Yasuhiro YOSHIDA*, Itushi UNO**, Takanori UCHIDA** and Yuji OHYA**.

* Department of Earth System Science and Technology, Kyushu University, Kasuga Park 6-1, Kasuga, Fukuoka, 816-8580, Japan.

** Research Institute for Applied Mechanics, Kyushu University.

(Received 16 August 2004 ; Accepted 15 September 2005)

日本気象学会英文レター誌 SOLA
(Scientific Online Letters on the Atmosphere)
<http://www.jstage.jst.go.jp/browse/sola/>

第1巻 2005年9月 目次

Masahiro HOSAKA, Daisuke NOHARA and Akio KITOH : Changes in Snow Cover and Snow Water Equivalent Due to Global Warming Simulated by a 20 km-mesh Global Atmospheric Model	93-96
Kazuo KURIHARA, Koji ISHIHARA, Hidetaka SASAKI, Yukio FUKUYAMA, Hitomi SAITO, Izuru TAKAYABU, Kazuyo MURAZAKI, Yasuo SATO, Seiji YUKIMOTO and Akira NODA : Projection of Climatic Change over Japan Due to Global Warming by High-Resolution Regional Climate Model in MRI	97-100
Kazuyo MURAZAKI, Hidetaka SASAKI, Hiroyuki TSUJINO, Izuru TAKAYABU, Yasuo SATO, Hiroshi ISHIZAKI and Kazuo KURIHARA : Climatic Change Projection for the Ocean around Japan Using a High-Resolution Coupled Atmosphere-Ocean Regional Climate Model	101-104
Masaru INATSU and Masahide KIMOTO : Difference of Boreal Summer Climate between Coupled and Atmosphere-only GCMs	105-108